



RESEARCH ARTICLE

# Effect of the Clinical Usage of Tetracaine Hydrochloride Ophthalmic Solution on Multiple Antibiotic Resistant *Staphylococcus Aureus*

Samuel Kyei<sup>1,2\*</sup>, Samuel Bert Boadi-Kusi<sup>1</sup>, Mary Nana Effriam-Menyah<sup>3</sup>, and Paulina Duah<sup>4</sup>

① Department of Optometry and Vision Science, School of Allied Health Sciences, University of Cape Coast, Ghana. | ② Biomedical and Clinical Research Centre, College of Health Allied Sciences, University of Cape Coast, Cape Coast. | ③ Eye Clinic, VRA Hospital, Akosombo, Ghana. | ④ Department of Medical Laboratory Science, School of Allied Sciences, University of Cape Coast, Cape Coast, Ghana.

\*Correspondence should be addressed to Samuel Kyei (email: skyei@ucc.edu.gh)

## Abstract

**Background:** The emergence of resistant strain bacteria is rendering many antibiotics ineffective in the management of infectious diseases.

**Objective:** The purpose was to investigate the effect of Tetracaine hydrochloride usage on the emergence of a resistant strain of bacteria.

**Materials and methods:** A total of fifteen New Zealand White rabbits of either sex weighing between 1400-2700 grams were used for this study. Infectious agents of Ophthalmia neonatorum were cultured, isolated and identified. Their susceptibility to various antibiotics was assessed. A resistant strain was picked up and inoculated on the conjunctiva of the rabbits to induce bacterial conjunctivitis. Rabbits were divided into three groups of five rabbits each. Group A received Tetracaine as treatment, group B received 0.3% Ciprofloxacin as treatment and group C received normal saline as a treatment for 14 days. Outcome measures included bacterial colony count, clinical signs and post-antibiotic susceptibility test.

**Results:** The tetracaine significantly reduced clinical signs ( $P=0.012$ ) compared to normal saline (control) and 0.3% Ciprofloxacin reduced clinical signs ( $P=0.003$ ) compared to normal saline (control). This means there was a significant decrease in clinical scores with the interventions as compared to the normal saline which showed minimal change in clinical score. Tetracaine hydrochloride significantly reduced bacteria load ( $P\leq 0.0001$ ). It decreased multiple antibiotic-resistant index by 20%.

**Conclusion:** Tetracaine has a supplemental antibiotic effect on resistant strain of bacteria and does not worsen antibiotic resistance of bacteria. Clinical usage of tetracaine ophthalmic solution should therefore be applied after conjunctival swaps or corneal scrapping have been taken.

**Keywords:** Antibiotics, Antimicrobial, *Staphylococcus aureus*, Tetracaine hydrochloride.

**Citation:** Kyei, S., Boadi-Kusi, S. B., Effriam-Menyah, M. N., Duah, P. (2023) Effect of the Clinical Usage of Tetracaine Hydrochloride Ophthalmic Solution on Multiple Antibiotic Resistant *Staphylococcus Aureus*. Integrated Health Research Journal 1(1), 1-6. <https://doi.org/10.47963/ihrij.v1i1.1173>

**Received** January 10, 2023; **Published** April 20, 2023

**Copyright:** ©2023 This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Introduction

Some studies have shown that anesthetics possess a supplemental antibiotic effect (Johnson, Saint John, & Dine, 2008). Several topical anesthetic agents have demonstrated distinct antimicrobial activity against specific bacterial strains and Candida (Pelosini, Treffene, & Hollick, 2009). Tetracaine hydrochloride ophthalmic solution is a local anesthetic indicated for processes requiring an ophthalmic anesthetic of fast and brief action.

Its usage for diagnostic purposes could promote bacterial resistance to antibiotics since they are used momentarily and not formulated for treatment. The possibility of repurposing older drugs in an attempt to deal with unmet medical needs further reinforces the necessity for such evaluation to inform the decision in this line of thought (Takahashi et al., 2017).

Evidence from research indicates the intrinsic antimicrobial characteristics of local anesthetics against

a broad spectrum of human pathogens (Johnson et al., 2008). At levels typically used in the clinical environment, multiple local anesthetics inhibit the development of numerous bacteria and fungi under different circumstances. Early research noted that Tetracaine exerts its bactericidal effect on the bacterial cell by a mechanism of action. The writers discovered that tetracaine harmed *Pseudomonas aeruginosa* cell membrane through lysis, intracellular leakage, dehydrogenase activity, and enhanced permeability of the cell wall and greater temperature correlates with a rise in the inhibition of microbial development. Observation of cell lysis, intracellular leakage, dehydrogenase activity, and greater sensitivity of spheroplasts to Tetracaine than whole cells led to the conclusion that Tetracaine works by damaging the cell membrane (Leung & Rawal, 1977). This study aimed to investigate the antibiotic effect of Tetracaine hydrochloride on a resistant strain of bacteria to ascertain its contribution to the increasing spate of resistance to conventional antibiotics.

## Materials and Methods

**Drugs and Chemical Used:** Tetracaine hydrochloride (Archdiocesan Health Pharmacy, Kumasi, Ghana) was used as an anesthetic and positive control drug was Ciprofloxacin eye drop (Ciron Drugs Ltd., India). Normal saline drops (MSB Pharma., India) was also used as a negative control drug in the study. Ketamine hydrochloride (Pfizer, USA) was used to anesthetize the rabbits before the induction of conjunctivitis.

**Animal husbandry:** A total of 15 New Zealand White rabbits of either sex weighing between 1.4kg-2.7kg were purchased from the School of Agriculture Rabbitry, University of Cape Coast and kept at the Animal House of the School of Biological Sciences of the University of Cape Coast, Ghana. The animals were housed singly in aluminum cages (34x47 x18 -cm<sup>3</sup>) with softwood shavings as bedding, under ambient laboratory conditions (28 ± 2°C, relative humidity 60%-70%, and a normal light-dark cycle) and fed with normal commercial pellet diet (AGRICARE, Kumasi) and water ad libitum.

**Culture and isolation of bacteria:** Swabs from clinically diagnosed cases of Ophthalmia Neonatorum were cultured, isolated and identified in the microbiology laboratory using standard microbiological procedures in a previous study (Kang & Lee, 1989). The susceptibility to various antibiotics was assessed using an antibiotic disc (Abtek Biologicals, Liverpool, UK). Multiple antibiotic-resistant strains of *Staphylococcus aureus* (*S. aureus*) as determined in a previous study (Kang & Lee, 1989) were picked from the pure culture with a sterile standard loop was inoculated into 10 ml of nutrient broth. The isolated multiple resistant *S. aureus* was then incubated for 24 hours at 37°C.

**Induction of infectious conjunctivitis:** The method as described by Al-Waili (Al-Waili, 2004) and adopted by Ilechie (Ilechie, Kwapong, Mate-Kole, Kyei, & Darko-Takyi, 2012) was used. Each rabbit was anesthetized by an intramuscular injection of 10mg/kg Ketamine

hydrochloride and a drop of 0.5% topical Tetracaine hydrochloride ophthalmic solution was instilled into the conjunctival cul-de-sac of the right eye of each animal. An abrasion was then made on the right bulbar conjunctiva of each rabbit. A 10µL of the bacterial suspension of multiple resistant *S. aureus* was inoculated into the abraded bulbar conjunctivae of the rabbits.

**Experimental design:** Rabbits were pre-examined for clinical signs of infectious conjunctivitis and those with no signs of conjunctivitis were selected and inoculated with the bacterial inoculum. Forty-eight hours after inoculation the rabbits that showed clinical signs of infectious conjunctivitis were selected and randomized for treatment (n=5). Swabs were taken to confirm positive cultures in the right eye of the animals before randomization.

They were randomized for treatment with 0.5% Tetracaine hydrochloride, 0.3% w/v of Ciprofloxacin (because it was the antibiotic that the strains were susceptible) and 0.9 w/v of Normal saline drop. All treatments were given twice daily (12 hourly intervals) for 14 days.

The right eyes of the animals were assessed for clinical signs of bacterial conjunctivitis using a handheld slit lamp every other day alongside swabbing for days of positive culture over the experimental period. The signs were scored and scale of 0-3 except for sensitivity to light. The scoring was as shown in **Table 1**.

The maximum possible score was 21. The swabs were used to assess for Bacteria Colony Count (BCC) and Antimicrobial Sensitivity Testing (AST). Plate count agar (PCA) was used for Bacteria Count Colony. The specimen swab was snapped off the swab shaft at the score line and cut shaft to fit into a test tube containing 9ml of sterilized freshly prepared peptone water. It was incubated at 37°C for 10h. Using a sterilized pipette, the culture was transferred into a sterilized tube containing 10ml of sterilized freshly prepared peptone water. Using a serial dilution method, 1ml of the aliquot was transferred into the sterilized tubes until least 10 x 5 dilution factor was reached. 1000µL of the serially diluted aliquot was transferred into an autoclaved media plate. Employing pour plate method, about 20ml of freshly prepared autoclaved media was then poured into the plates, swirled gently in both clockwise and anti-clockwise directions and allowed to solidify. The solidified plates were labelled and incubated (Panasonic MIR-154-PE Cooled incubator) at 37 °C for 24 hours. Samples in VRBA were allowed a maximum incubation time of 1 week. Two different dilution factors for each sample was used and the mean found. After incubation, single viable colonies were counted and recorded in CFU/ml, using the formula; CFU/mL = CFU x dilution factor x 1/aliquot.

Modified Kirby-Bauer disc diffusion was employed. 8ml of normal saline was inoculated with portions of at least 3 pure colonies with similar appearance using a hot-flamed sterilized straight wire. The turbidity of the suspension was then compared with 0.5 McFarland turbidity standard (1.5x10<sup>8</sup>CFU/ml) and adjusted by either the addition of more colonies or normal saline. Freshly prepared Mueller Hinton (MH) agar in sterilized media plates were streaked with a sterilized swab stick soaked with the inoculums in

Table 1: Clinical and visual scoring of some features associated with bacterial conjunctivitis

Clinical features	0	1 (mild)	2 (moderate)	3 (severe)
Redness	Clear	Redness size of <2mm	Redness size of 2-5mm	Redness of >5mm
Discharge	None	Mild discharge	Moderate discharge	Severe discharge
Crusty eyelids	None	Mild crusty eyelids	Moderate crusty eyelids	Severe crusty eyelids
Wet eye	None	Mild wet eye	Moderate wet eye	Severe wet eyes
Edema	No edema present in the conjunctiva	Slight edema/general or localized conjunctival edema	General diffuse edema of the conjunctiva	Frank edema of chemosis of conjunctiva
Photophobia	No photophobia	slight difficulty with light causing occasional eye	Moderate difficulty with light requiring regular blinking	Severe difficulty with light cannot bear natural light
Blepharospasm	No spasm	Mild spasm at stimulation only	Visible spasm without impairment of daily life	Visible spasm with impairment of daily life
Photophobia	No photophobia	slight difficulty with light causing occasional eye	Moderate difficulty with light requiring regular blinking	Severe difficulty with light cannot bear natural light
Blepharospasm	No spasm	Mild spasm at stimulation only	Visible spasm without impairment of daily life	Visible spasm with impairment of daily life

the normal saline suspension. The whole surfaces of the media were streaked three (3) times at an angle 60°. The plates were rotated each time, and allowed to air dry. Sterilized hot-flame forceps were used to place the gram appropriate antimicrobial-impregnated disk (Axiom) on the agar, and each disc was pressed gently to ensure complete contact with the surface of the agar. The plates were then incubated at 37 °C for 24 hours. After 24 hours, the diameters of inhibition zones were measured using a rule and recorded in millimeters (mm). Antimicrobial sensitivity pattern was determined as susceptible (S), intermediate (I) or resistant (R), with reference to AST interpretation chart.

*Statistical analysis:* Data obtained for control, test and reference drug effects were analyzed by one-way analysis of variance followed by Tukey’s multiple comparisons test using GraphPad Prism (version 5.03; GraphPad, La Jolla, CA, USA). Values were expressed as the mean ± standard error of the mean. P≤0.05 was considered to be statistically significant.

## Results

After 48 hours of inoculation of *S. aureus*, there were obvious signs of bacterial conjunctivitis in all treatment groups. The rabbits treated with normal saline showed significantly higher clinical signs of bacterial conjunctivitis as compared to the 0.3% Ciprofloxacin and Tetracaine hydrochloride treatment groups (Table 2).

There was a significant reduction in redness, discharge, crusty eyelids and conjunctiva edema relative to the control; seen as significant decrements (P≤0.001-0.004) in clinical scores of bacteria conjunctivitis (Table 2, Figure 1). However, tetracaine significantly reduced clinical signs (P=0.012) compared to normal saline (control) and 0.3% Ciprofloxacin reduced clinical signs (P =0.003) compared to normal saline (control). This means there was a significant decrease in clinical scores with the interventions as compared to the normal saline which showed minimal change in clinical score (Figure 1, Table 2).

Table 2: The effect of Normal saline (control), Ciprofloxacin and Tetracaine on clinical scores per treatment schedule

Day	Clinical Scores		
	Normal Saline	Ciprofloxacin	Tetracaine
1	13.00±2.12	10.8±0.530	11.4±2.088
3	9.200±2.332	4.4±1.030	6.8±0.860
5	8.400±0.812	5.4±0.40	7.8±0.735
7	8.400±0.510	4.8±0.374*	3.8±0.735*
9	7.600±0.872	4.8±0.374	3.8±0.490*
11	8.600±0.927	3.8±0.735***	4.8±0.970
13	8.400±0.600	4.6±0.675**	3.6±0.400***
15	6.800±0.860*	4.2±0.490***	2.8±0.490***

Significant clinical score on treatment days established using a one-way analysis of variance (ANOVA) followed by Tukey’s multiple comparisons test. \* P≤0.05; \*\*P≤0.01, P≤0.001.




Figure 1: The effect of normal saline (control), 0.3% ciprofloxacin and tetracaine on clinical scores. Significant clinical score per treatment established using one-way analysis of variance (ANOVA) followed by Tukey’s multiple comparisons test. \* P=0.012, \*\*P=0.003.

*Bacteria colony count:* There was a significant decrease (P=0.002) in the bacteria colony count of rabbits treated with the Ciprofloxacin as compared to the control group (treated with normal saline) (Figure 2). Tetracaine hydrochloride exhibited a significant (P=0.004) antibiotic effect on the multiple antibiotic-resistant *S. aureus* (Table 3).




Figure 2: The effect of Normal saline (control), 0.3% Ciprofloxacin and Tetracaine on bacteria count after treatment. Significant reductions in bacteria count per treatment established using a one-way analysis of variance (ANOVA) followed by Tukey's multiple comparisons test. \*\*implies  $P \leq 0.002-0.004$ .

Table 3: The effect of Normal saline (control), 0.3% Ciprofloxacin and Tetracaine on bacteria count per treatment schedule.

Day	Bacteria Count		
	Normal Saline	Ciprofloxacin	Tetracaine
1	22348±8749.50	19074.0±6279.15	18945.80±4411.30
3	23632±11391.15	4532.00±1441.023	3262.80±484.85
5	10406±3989.39	2766.00±688.56*	2718.00±199.16*
7	8944±3542.020	2542.20±694.61*	2177.00±302.29*
9	6694±29739.86	1415.00±950.65*	2087.20±387.39*
11	4828±100.10	1058.60±824.00*	1276.40±379.13*
13	4208±900.80	920.00±676.246*	617.40±283.05*
15	3168±463.41	337.80±173.10**	375.60±153.93**

Significant reductions in bacteria count per treatment days established using a one-way analysis of variance (ANOVA) followed by Tukey's multiple comparisons test. \* $P=0.002$ ; \*\* $P=0.004$

**Multiple antibiotic resistant index:** The antibiotic susceptibility test did not show any significant multiple antibiotic-resistant in bacteria isolates treated with Tetracaine hydrochloride and normal saline (control). However, there was a significant increase in multiple antibiotic-resistance in isolates treated with Ciprofloxacin (Figure3; Table 4).




Figure 3: The MAR Indexes for all treatment groups. Significant reduction in MAR Index of treatment groups established using a one-way analysis of variance (ANOVA) followed by Tukey's multiple comparisons test. \*\*implies  $P \leq 0.01$ .

Table 4: The MAR Indexes for Normal saline, Ciprofloxacin, and Tetracaine treatment groups

Day	MAR Index values (%)		
	Normal Saline	Ciprofloxacin	Tetracaine
1	25	62.5	12.5
3	37.5	50	25
5	25	62.5	25
7	25	37.5	12.5
9	25	75	12.5
Average	27.500±2.500	57.500±6.374 **	17.500±3.062

One-way analysis of variance (ANOVA) followed by Tukey's multiple comparisons test. \*\*implies  $P \leq 0.01$ . MAR, Multiple antibiotic resistance define MAR here

## Discussion

Bacterial conjunctivitis can be contracted directly from infected individuals or can result from abnormal proliferation of the native conjunctival flora (Azari & Barney, 2013). Contaminated fingers (Høvding, 2008), oculogenital spread (Varu et al., 2019) and contaminated fomites (Sattar, Dimock, Ansari, & Springthorpe, 1988) are common routes of transmission. Besides, certain conditions such as compromised tear production, disruption of the natural epithelial barrier, abnormality of adnexal structures, trauma, and immune-suppressed status predispose individuals to bacterial conjunctivitis (Varu et al., 2019). The most common pathogens for bacterial conjunctivitis in adults are Staphylococcal species, followed by *Streptococcus pneumoniae* and *Haemophilus Influenza* (Fishovitz, Hermoso, Chang, & Mobashery, n.d.). In children, the disease is often caused by *Haemophilus influenzae*, *Streptococcus pneumoniae* and *Moraxella catarrhalis* (Fishovitz et al., n.d.). The course of the disease usually lasts 7 to 10 days the reason for the decline in the clinical scores of conjunctivitis in the rabbits in the control group (Boadi-Kusi et al., 2021). Signs and symptoms include red eye, purulent or mucopurulent discharge, and chemosis (Fishovitz et al., n.d.). The period of incubation and communicability is estimated to be 1 to 7 days and 2 to 7 days, respectively (Leung & Rawal, 1977). Bilateral matting of the eyelids and adherence of the eyelids, lack of itching, and no history of conjunctivitis were strong positive predictors of bacterial conjunctivitis (Sharma, 2011). This study aimed to investigate the antibiotic effect of Tetracaine hydrochloride on a resistant strain of bacteria to ascertain its contribution to the increasing spate of resistance to conventional antibiotics.

In this study resistant strain of *Staphylococcus aureus* was the causative agent of bacterial conjunctivitis. There was a significant reduction in clinical signs of bacterial conjunctivitis in the rabbits that were treated with Tetracaine hydrochloride ophthalmic solution. An earlier study (Reynolds, Greenwood-Quaintance, Patel, & Pulido, 2016) reported on the mechanism of action by which Tetracaine damaged the cell wall of bacteria through lysis, leakage of intracellular components, dehydrogenase activity and increased cell wall permeability.

A study on the effects of topical anesthetics on bacteria using a disk diffusion technique studied proparacaine and tetracaine at 0.5, 0.25, and 0.125% concentrations

found that tetracaine inhibited *S. aureus* growth at 5000 µg/mL and *Pseudomonas aeruginosa* at 2500 to 5000 µg/mL (Chiang, Penadés, & Chen, 2019). Also, a study using broth microdilution demonstrated that tetracaine inhibited strains of *S. epidermidis* at a concentration of 625µg/mL (Mullin & Rubinfeld, 1997). Tetracaine has demonstrated its antibiotic effects against coagulase-negative staphylococci and *S. aureus* (Johnson et al., 2008).

In the present study, bacterial colonies were counted from bacterial-induced conjunctivitis in rabbit eye treated with tetracaine and there was a significant reduction in bacteria count. This particular solution contained 0.5% Tetracaine hydrochloride as an active ingredient, a preservative which is Chlorobutanol and Boric Acid, Edetate Disodium, Potassium Chloride as inactive ingredients.

Bacteria may have innate resistance or acquired resistance to antimicrobial agents. This involves; mutations in cell genes (chromosomal mutation) leading to resistance, transformational gene transfer from one microorganism to another, conjugation which needs independent genetic elements including transposons (Tns) and plasmids and also transduction which involves independently replicating bacterial viruses, known as bacteriophages (Kaewjiaranai, Srisatjaluk, Sakdajeyont, Pairuchvej, & Wongsirichat, 2018). The current class of antimicrobial agents targets certain major bacterial processes such as metabolic pathways, cell wall, functions of the cell membrane, protein and nucleic acid synthesis (Rietveld, van Weert, ter Riet, & Bindels, 2003). After the bacteria have gained resistance genes against various antimicrobial agents, the bacteria can, therefore, use several biochemical types of resistance mechanisms including drug inactivation or enzymatic degradation involving enzymes that destroy the antibiotic. These enzymes include B-lactamases that cleave the amide bond in the B-lactam ring (Loka, Sumadja, & Resmi, 2017). Another mechanism is by extrusion of the antibiotic before it reaches the target by the efflux pump. This mechanism extrudes the antibiotic before it reaches the bacteria target-site (Kaewjiaranai et al., 2018). Target modification occurs by an altered bacteria cell wall which modifies the target region. An example is MRSA which involves a modified Penicillin Binding Protein (PBP2a) that confers methicillin resistance to certain *S. aureus* strains (Loka et al., 2017).

In this study, the antibiotic susceptibility test on bacteria isolates treated with 0.3% Ciprofloxacin showed multiple antibiotic-resistant which was not the case before treatment was commenced. There was no increase in multiple antibiotics resistant in groups treated with Tetracaine hydrochloride and normal saline. A study by Sharma (Sharma, 2011), among 2408 eyes in South Florida revealed an increase resistance of gram-positive by 2-fold and 3-fold to erythromycin and ciprofloxacin respectively. The primary disadvantage associated with treatment with an antibiotic is the probable future resistance. The findings of this study support the assertion that there exists a link between the consumption of antibiotics and subsequent resistance (Leibovici et al., 2001). This was not the case Tetracaine hydrochloride treatment given credence to the fact that the short to medium term diagnostic does not

contribute to the growing spate of antibiotic resistance. The limitation to this study was that there were no genetic studies to confirm this finding.

## Conclusion

Despite Tetracaine possessing supplemental antibiotic effect as a diagnostic agent, it has no consequential effect on the development of antibiotic-resistant strains of bacteria. It is recommended that further studies involving molecular methods be adopted to confirm this finding.

## Declarations

*Ethical considerations:* The study protocol was approved by the University of Cape Coast Institutional Review Board, UCCIRB, Ghana with reference number: UCCIRB/CHAS/2018/57. All activities performed during the studies conformed to the Association for Research in Vision and Ophthalmology Statement for Use of Animals in Ophthalmic and Vision Research. Biosafety guidelines for the protection of personnel in the laboratory were observed.

*Availability of data and materials:* All data generated or analysed during this study are included in this published article.

*Authors' contributions:* Author SK conceived the idea, designed the study and wrote the protocol. SBB-K managed the analyses and the interpretation of data. ME-M managed the literature searches, collected data, and wrote the first draft of the manuscript. All authors read, critically revised the content and approved the final manuscript.

*Acknowledgment:* The authors are grateful to Esther Duncan for editing and formatting the manuscript.

*Conflict of Interest:* The authors declare no conflict of interest.

## References

- Al-Waili, N. S. (2004). Investigating the antimicrobial activity of natural honey and its effects on the pathogenic bacterial infections of surgical wounds and conjunctiva. *Journal of Medicinal Food*, 7(2), 210–222.
- Azari, A. A., & Barney, N. P. (2013). Conjunctivitis: a systematic review of diagnosis and treatment. *Jama*, 310(16), 1721–1730.
- Boadi-Kusi, S. B., Kyei, S., Holdbrook, S., Abu, E. K., Ntow, J., & Ateko, A. M. (2021). A study of Ophthalmia Neonatorum in the Central Region of Ghana: Causative Agents and Antibiotic Susceptibility Patterns. *Global Pediatric Health*, 8, 2333794X211019700.
- Chiang, Y. N., Penadés, J. R., & Chen, J. (2019). Genetic transduction by phages and chromosomal islands: The new and noncanonical. *PLoS Pathogens*, 15(8), e1007878.
- Fishovitz, J., Hermoso, J. A., Chang, M., & Mobashery, S. (n.d.). Critical Review Penicillin-binding Protein 2a of.
- Høvding, G. (2008). Acute bacterial conjunctivitis. *Acta Ophthalmologica*, 86(1), 5–17.
- Ilechie, A. A., Kwapong, P. K., Mate-Kole, E., Kyei, S., & Darko-

- Takji, C. (2012). The efficacy of stingless bee honey for the treatment of bacteria-induced conjunctivitis in guinea pigs. *Journal of Experimental Pharmacology*, 63–68.
- Johnson, S. M., Saint John, B. E., & Dine, A. P. (2008). Local anesthetics as antimicrobial agents: a review. *Surgical Infections*, 9(2), 205–213.
- Kaewjiaranai, T., Srisatjaluk, R. L., Sakdajeyont, W., Pairuchvej, V., & Wongsirichat, N. (2018). The efficiency of topical anesthetics as antimicrobial agents: A review of use in dentistry. *Journal of Dental Anesthesia and Pain Medicine*, 18(4), 223–233.
- Kang, S. W., & Lee, J. H. (1989). Antibacterial effect of tetracaine hydrochloride (Pontocaine (R)) in the aspect of exposure time: in vitro study. *Korean Journal of Ophthalmology*, 3(1), 11–13.
- Leibovici, L., Berger, R., Gruenewald, T., Yahav, J., Yehezkeili, Y., Milo, G., ... Pitlik, S. D. (2001). Departmental consumption of antibiotic drugs and subsequent resistance: a quantitative link. *Journal of Antimicrobial Chemotherapy*, 48(4), 535–540.
- Leung, Y.-W., & Rawal, B. D. (1977). Mechanism of action of tetracaine hydrochloride against *Pseudomonas aeruginosa*. *Journal of Infectious Diseases*, 136(5), 679–683.
- Loka, W. P., Sumadja, W. A., & Resmi. (2017). *Journal of Chemical Information and Modeling*, 21(2), 1689–1699. Retrieved from <https://www.oecd.org/dac/accountable-effective-institutions/Governance-Notebook-2.6-Smoke.pdf>
- Mullin, G. S., & Rubinfeld, R. S. (1997). The antibacterial activity of topical anesthetics. *Cornea*, 16(6), 662–665.
- Pelosini, L., Treffene, S., & Hollick, E. J. (2009). Antibacterial activity of preservative-free topical anesthetic drops in current use in ophthalmology departments. *Cornea*, 28(1), 58–61.
- Reynolds, M. M., Greenwood-Quaintance, K. E., Patel, R., & Pulido, J. S. (2016). Selected antimicrobial activity of topical ophthalmic anesthetics. *Translational Vision Science & Technology*, 5(4), 2.
- Rietveld, R. P., van Weert, H. C. P. M., ter Riet, G., & Bindels, P. J. E. (2003). Diagnostic impact of signs and symptoms in acute infectious conjunctivitis: systematic literature search. *Bmj*, 327(7418), 789.
- Sattar, S. A., Dimock, K. D., Ansari, S. A., & Springthorpe, V. S. (1988). Spread of acute hemorrhagic conjunctivitis due to enterovirus-70: Effect of air temperature and relative humidity on virus survival on fomites. *Journal of Medical Virology*, 25(3), 289–296.
- Sharma, S. (2011). Antibiotic resistance in ocular bacterial pathogens. *Indian Journal of Medical Microbiology*, 29(3), 218–222.
- Takahashi, K., Wang, F., Kantarjian, H., Doss, D., Khanna, K., Thompson, E., ... Gumbs, C. (2017). Preleukaemic clonal haemopoiesis and risk of therapy-related myeloid neoplasms: a case-control study. *The Lancet Oncology*, 18(1), 100–111.
- Varu, D. M., Rhee, M. K., Akpek, E. K., Amescua, G., Farid, M., Garcia-Ferrer, F. J., ... Dunn, S. P. (2019). Conjunctivitis preferred practice pattern®. *Ophthalmology*, 126(1), P94–P169.